

Techniques and Procedures

REMOVAL OF THE PREHOSPITAL TOURNIQUET IN THE EMERGENCY DEPARTMENT

Matthew J. Levy, DO, MSc,* Jason Pasley, DO,† Kyle N. Remick, MD,‡ Alexander L. Eastman, MD, MPH,§
Asa M. Margolis, DO, MPH, MS,* Nelson Tang, MD,* and Craig A. Goolsby, MD, MED‡

*Department of Emergency Medicine, School of Medicine, Johns Hopkins University, Baltimore, Maryland, †Department of Surgery, School of Medicine, Michigan State University, East Lansing, Michigan, ‡McLaren Oakland Hospital Pontiac, Michigan Uniformed Services University of the Health Sciences, Bethesda, Maryland, and §U.S. Department of Homeland Security, Washington, DC

Reprint Address: Matthew J. Levy, DO, MSc, Department of Emergency Medicine, School of Medicine, Johns Hopkins University, 5801 Smith Ave., Davis Building Suite 3200, Baltimore, MD 21209

Abstract—Background: Life-threatening hemorrhage from extremity injuries can be effectively controlled in the prehospital environment through direct pressure, wound packing, and the use of tourniquets. Early tourniquet application has been prioritized for rapid control of severe extremity hemorrhage and is a cornerstone of prehospital trauma resuscitation guidelines. Emergency physicians must be knowledgeable regarding the initial assessment and appropriate management of patients who present with a prehospital tourniquet in place. **Discussion:** An interdisciplinary group of experts including emergency physicians, trauma surgeons, and tactical and Emergency Medical Services physicians collaborated to develop a stepwise approach to the assessment and removal (discontinuation) of an extremity tourniquet in the emergency department after being placed in the prehospital setting. We have developed a best-practices guideline to serve as a resource to aid the emergency physician in how to safely remove a tourniquet. The guideline contains five steps that include: 1) Determine how long the tourniquet has been in place; 2) Evaluate for contraindications to tourniquet removal; 3) Prepare for

tourniquet removal; 4) Release the tourniquet; and 5) Monitor and reassess the patient. **Conclusion:** These steps outlined will help emergency medicine clinicians appropriately evaluate and manage patients presenting with tourniquets in place. Tourniquet removal should be performed in a systematic manner with plans in place to immediately address complications. © 2020 Elsevier Inc. All rights reserved.

Keywords—stop the Bleed; prehospital tourniquet; tourniquet removal; tourniquet conversion

INTRODUCTION

Hemorrhage accounts for 35% of prehospital trauma deaths and nearly 40% of deaths within the first 24 h of injury (1). Life-threatening bleeding from extremity injuries can be effectively controlled in the prehospital environment through direct pressure, wound packing, and the use of tourniquets. Tourniquet application has been shown to be safe, particularly if the tourniquet can be removed within 2 h of application (2–4). Placement of prehospital tourniquets in the civilian population has also been associated with a sixfold survival benefit (5). Early tourniquet application has been prioritized for rapid control of life-threatening extremity hemorrhage and is a cornerstone of prehospital trauma resuscitation

Disclaimer: The views expressed in this manuscript reflect those of the authors alone and do not reflect the official policy or position of the Uniformed Services University of the Health Sciences, Department of the Army, Department of the Air Force, Department of Defense, or the United States Government.

guidelines (6–9). Tourniquets are increasingly being carried by law enforcement officers, first responders, and Emergency Medical Services (EMS) units (10,11). Studies have shown that laypersons can properly apply tourniquets with minimal training (12,13). Programs, including Stop the Bleed, empower laypeople with knowledge and skills, including tourniquets, to stop life-threatening bleeding (14–17). Tourniquets may be applied by members of the general public, first responders, and EMS clinicians to patients suffering from life-threatening extremity hemorrhage.

Emergency physicians must be knowledgeable regarding the initial assessment and appropriate management of patients who present with a tourniquet in place. This includes the indications, contraindications, proper methods, and potential pitfalls for tourniquet removal or conversion to another bleeding control maneuver or intervention. All efforts should be made to convert tourniquets to another type of bleeding control mechanism as soon as it can be done safely, and within 2 h when at all possible (7,8,18–20). Tourniquets in place for more than 2 h, especially those in place 6 h or longer, should be removed in a critical care setting, with the capability to address the local and systemic reperfusion effects associated with prolonged ischemia, as well as treat rhabdomyolysis, compartment syndrome, and other possible complications (21,22).

DISCUSSION

An interdisciplinary group of experts including emergency physicians, trauma surgeons, and tactical and EMS physicians collaborated to develop a stepwise approach to the assessment and removal of a limb tourniquet in the emergency department (ED) after being placed in the prehospital setting. All members of this group were established subject matter experts and have credibility in multiple disciplines directly related to management of life-threatening hemorrhage including: emergency medicine (MJL, AMM, NT, CAG), trauma surgery (JP, KR, ALE), military medicine (JP, KR, CAG), and emergency medical services medicine (MJL, AMM, ALE, NT). In addition, multiple authors maintain national leadership roles within the Stop the Bleed community in the United States (MJL, ALM, CAG). The goal of this group was to share a best-practices approach to serve as a simple, yet comprehensive, resource for emergency physicians and resuscitation team members. A search of the existing literature was conducted and served as the base for the formation of the technique presented. This technique assumes familiarity with evaluation of the injured patient using standard best practices for initial trauma and critical care resuscitation.

Technique: See Figure 1 (flow diagram)

1. *Determine how long the tourniquet has been in place and the availability of surgical resources.* If the exact duration cannot be determined, use time of EMS arrival to the scene.
 - a. If the tourniquet has been in place < 90 min, and surgical care is immediately available (within 30 min of patient arrival in the ED), immediately obtain a surgical consult. This is to help minimize the duration of tourniquet time to < 120 min and thus, help minimize ischemic tissue injury.
 - b. If the tourniquet has been in place < 90 min and surgical support is not immediately available, determine if the patient can be transferred to, and arrive at, a trauma center within < 120 min of tourniquet placement. If so, transfer to trauma center.
 - i. If the tourniquet is, or will be, in place > 120 min, consider consultation with the trauma surgeon at the trauma center, and assess the individual patient's circumstances (including presence of shock, other life-threatening emergencies, and the severity of the wound) and evaluate the risk of ischemia compared with the risk of re-bleeding secondary to tourniquet removal.
 - c. If a tourniquet has been in place > 120 min, assess for a contraindication to tourniquet removal.
 - i. If the treating physician determines that it is in the patient's best interests to remove a tourniquet that has been in place for ≥ 2 h, we highly encourage this be done in a critical care setting (ED/intensive care unit/operating room) that has resources including monitoring, fluids, resuscitation medications, laboratory, and surgical capability.
2. *Evaluate for Contraindications to Tourniquet Removal:* The following conditions are considered contraindications to tourniquet removal.
 - a. Traumatic amputation with a tourniquet placed immediately proximal to the amputation stump (within several inches).
 - b. Decompensated shock/other life-threatening injuries/impending cardiopulmonary arrest.
 - i. After the patient is stabilized, the tourniquet may be evaluated for removal as part of the secondary survey.
 - c. Inability to closely monitor wound for signs of rebleeding.
 - i. Once a tourniquet is removed, direct patient monitoring is required for a minimum of 1 h for signs of significant re-bleeding.




Figure 1. Emergency department approach to the patient with a prehospital tourniquet.

3. *Prepare for Tourniquet Removal:* If no contraindications to tourniquet removal are present, and sufficient resources exist, consider having an additional new tourniquet immediately available in the event that reapplication of the tourniquet becomes necessary and the initial tourniquet is damaged. Additional equipment, including hemostatic gauze, dressing materials, cardiac monitoring, and Advanced Cardiac Life Support medications should also be readily available.

- a. Clinicians should be aware that tourniquets are subject to significant wear and tear during application, therefore, having access to a new tourniquet, if at all possible, is advised (23–25).
- b. If the initial tourniquet was placed very high (proximally), a second tourniquet may be prepositioned in a location closer to the injury (within a few inches) but not directly over a joint.

4. Release the Tension on the Tourniquet:
 - a. If life-threatening bleeding is noted from the wound, re-tighten the tourniquet (or, if necessary, apply the second/backup tourniquet), noting the time of tourniquet reapplication. Prepare the patient for emergency surgery or transfer to a trauma center.
 - b. If no active bleeding, apply a dressing per routine practice.
 - i. Continue to monitor the wound for signs of recurrent bleeding.
 - c. If non-life-threatening bleeding is present:
 - i. A hemostatic agent-impregnated gauze (preferred, if available) should be packed into the wound.
 1. Manual pressure should be held for a minimum of 3–5 min.
 2. Firmly secure the gauze in place with a pressure dressing.
 - ii. If no hemostatic gauze is available, sterile gauze should be used to pack into the wound.
 1. Apply manual pressure followed by a firmly secured pressure dressing after bleeding has stopped.
 - iii. If blood rapidly soaks the recently placed packing, tourniquet removal (conversion) has failed.
 1. Tighten the tourniquet within a few inches of the wound, but not over a joint.
 2. If bleeding still persists, apply a second tourniquet proximal to the first tourniquet. Make arrangements for urgent surgical evaluation or transfer to a trauma center.
 - d. After successful conversion of the tourniquet to a pressure dressing, assess the extremity for the presence of distal pulses, and motor and sensory function.
 - i. If no further bleeding is noted, keep the tourniquet loose on the limb, in case needed for re-bleeding.
5. *Monitor & Reassess*: Once a patient with a tourniquet has been converted successfully to a conventional dressing, it is essential that the patient undergo serial reassessments for evidence of re-bleeding and to ensure the presence of distal neurovascular function.
 - a. Continue diagnostic evaluation and management, while carefully observing the patient for recurrence of bleeding for a minimum of 1 h.
 - b. As the patient's condition stabilizes and blood pressure increases, so do the chances for rebleeding.
 - i. The previously placed tourniquet(s) should remain loose and proximal to the injury in case prompt reapplication is needed.
 - c. If reapplication is needed, reassessment of the patient and their wound is indicated, and definitive surgical control may be needed.

CONCLUSION

Life-threatening extremity bleeding may occur after instances of everyday traumatic injuries, as well as in mass casualty and high-threat situations. Patients will arrive at EDs via EMS and also via self-presentation, with tourniquets in place. These steps outlined will help emergency medicine clinicians initially caring for the traumatically injured patient to appropriately evaluate and manage patients presenting with tourniquets in place. Tourniquet removal (conversion) in the ED should be performed in a systematic manner, with plans in place to address complications.

REFERENCES

1. Kauvar DS, Lefering R, Wade CE. Impact of hemorrhage on trauma outcome: an overview of epidemiology, clinical presentations, and therapeutic considerations. *J Trauma* 2006;60(6 suppl):S3–11.
2. Kragh JF Jr, Walters TJ, Baer DG, et al. Practical use of emergency tourniquets to stop bleeding in major limb trauma. *J Trauma* 2008; 64(2 suppl):S38–50.
3. Wakai A, Winter DC, Street JT, Redmond PH. Pneumatic tourniquets in extremity surgery. *J Am Acad Orthop Surg* 2001;9:345–51.
4. Hagenouw RR, Bridenbaugh PO, van Egmond J, Stuebing R. Tourniquet pain: a volunteer study. *Anesth Analg* 1986;65:1175–80.
5. Teixeira PGR, Brown CVR, Emigh B, et al. Civilian prehospital tourniquet use is associated with improved survival in patients with peripheral vascular injury. *J Am Coll Surg* 2018;226:769–7761.
6. Jacobs LM, McSwain NE Jr, Rotondo MF, et al. Improving survival from active shooter events: the Hartford Consensus. *J Trauma Acute Care Surg* 2013;74:1399–400.
7. National Association of Emergency Medical Technicians Tactical Combat Casualty Care – Medical Personnel (NAEMT TCCC-MP) guidelines and curriculum. Available at: https://www.naemt.org/docs/default-source/education-documents/tccc/tccc-mp-updates-190801/tccc-guidelines-for-medical-personnel-190801.pdf?sfvrsn=cc99d692_2. Accessed September 22, 2022.
8. Committee for Tactical Emergency Care (C-TECC) Guidelines. Available at: <http://www.c-tecc.org/guidelines/als-blis>. Accessed September 22, 2020.
9. Rasmussen TE, Baer DG, Goolsby C. The giving back: battlefield lesson to national preparedness. *J Trauma Acute Care Surg* 2016; 80:166–7.
10. Doyle GS, Taillac PP. Tourniquets: a review of current use with proposals for expanded prehospital use. *Prehosp Emerg Care* 2008;12: 241–56.
11. Goodwin T, Moore KN, Pasley JD, Troncoso R Jr, Levy MJ, Goolsby C. From the battlefield to main street: tourniquet acceptance, use, and translation from the military to civilian settings. *J Trauma Acute Care Surg* 2019;87(1S suppl 1):S35–9.
12. Goolsby C, Branting A, Chen E, Mack E, Olsen C. Just-in-time to save lives: a pilot study of layperson tourniquet application. *Acad Emerg Med* 2015;22:1113–7.

13. Goolsby C, Chen E, Branting A, et al. Analysis of layperson tourniquet application using a novel color-coded device. *Disaster Med Public Health Prep* 2016;10:274–80.
14. The White House. FACT SHEET: Bystander: “Stop the Bleed” broad private sector support for effort to save lives and build resilience. Available at: <https://obamawhitehouse.archives.gov/the-press-office/2015/10/06/fact-sheet-bystander-stop-bleed-broad-private-sector-support-effort-save>. Accessed September 22, 2020.
15. Goolsby C, Jacobs L, Hunt RC, et al. Stop the Bleed Education Consortium: education program content and delivery recommendations. *J Trauma Acute Care Surg* 2018;84:205–10.
16. Jacobs LM Jr. Joint Committee to Create a National Policy to Enhance Survivability from Intentional Mass-Casualty and Active Shooter Events. The Hartford Consensus III: implementation of bleeding control—if you see something do something. *Bull Am Coll Surg* 2015;100:20–6.
17. Levy MJ, Jacobs LM. A call to action to develop programs for bystanders to control severe bleeding. *JAMA Surg* 2016;151:1103–4.
18. Dayan L, Zinmann C, Stahl S, Norman D. Complications associated with prolonged tourniquet application on the battlefield. *Mil Med* 2008;173:63–6.
19. Rorabeck CH. Tourniquet-induced nerve ischemia: an experimental investigation. *J Trauma* 1980;20:280–6.
20. Kragh JF Jr, O’Neill ML, Walters TJ, et al. Minor morbidity with emergency tourniquet use to stop bleeding in severe limb trauma: research, history, and reconciling advocates and abolitionists. *Mil Med* 2011;176:817–23.
21. Shackelford SA, Butler FK Jr, Kragh JF Jr, et al. Optimizing the use of limb tourniquets in Tactical Combat Casualty Care: TCCC Guidelines Change 14-02. *J Spec Oper Med* 2015;15:17–31.
22. Walters TJ, Mabry RL. Issues related to the use of tourniquets on the battlefield. *Mil Med* 2005;170:770–5.
23. Childers R, Tolentino JC, Leasiolagi J, et al. Tourniquets exposed to the Afghanistan combat environment have decreased efficacy and increased breakage compared to unexposed tourniquets. *Mil Med* 2011;176:1400–3.
24. Weppner J, Lang M, Sunday R, Debiase N. Efficacy of tourniquets exposed to the afghanistan combat environment stored in individual first aid kits versus on the exterior of plate carriers. *Mil Med* 2013;178:334–7.
25. Kragh JF Jr, O’Neill ML, Walters TJ, et al. The military emergency tourniquet program’s lessons learned with devices and designs. *Mil Med* 2011;176:1144–52.